PATIENT HEALTH QUESTIONNAIRE (PHQ-9)

NAME:	DATE:			
Over the last 2 weeks, how often have you been	Date o	of Bir	ths	
bothered by any of the following problems? (use "✓" to indicate your answer)	Not at all	Several days	More than half the days	Nearly every day
Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	О	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself—or that you are a failure or have let yourself or your family down	O	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed. Or the opposite — being so figety or restless that you have been moving around a lot more than usual.	0	1	2	3
9. Thoughts that you would be better off dead, or of hurting yourself	0	1	2	3
	add colum	ns	÷	÷
(Healthcare professional: For interpretation of TC please refer to accompanying scoring card).	TAL. TOTA	\L:		
10. If you checked off any problems, how difficult	Not difficult at all			
have these problems made it for you to do	Somewhat difficult			it
your work, take care of things at home, or get	Very difficult			
along with other people?				

Copyright € 1999 Pfizer Inc. All rights reserved. Reproduced with permission PRIME-MD€ is a trademark of Pfizer Inc. A2663B 10-C4-2005

Extremely difficult

PHQ-9 Patient Depression Questionnaire

For initial diagnosis:

- 1. Patient completes PHQ-9 Quick Depression Assessment.
- 2. If there are at least 4 √s in the shaded section (including Questions ±1 and ±2), consider a depressive disorder. Add score to determine severity.

Consider Major Depressive Disorder

- if there are at least 5 ✓s in the shaded section (one of which corresponds to Question #1 or #2)

Consider Other Depressive Disorder

- if there are 2-4 √s in the shaded section (one of which corresponds to Question #1 or #2)

Note: Since the questionnaire relies on patient self-report, all responses should be verified by the clinician, and a definitive diagnosis is made on clinical grounds taking into account how well the patient understood the questionnaire, as well as other relevant information from the patient.

Diagnoses of Major Depressive Disorder or Other Depressive Disorder also require impairment of social, occupational, or other important areas of functioning (Question #10) and ruling out normal bereavement, a history of a Manic Episode (Bipolar Disorder), and a physical disorder, medication, or other drug as the biological cause of the depressive symptoms.

To monitor severity over time for newly diagnosed patients or patients in current treatment for depression:

- Patients may complete questionnaires at baseline and at regular intervals (eg. every 2 weeks) at home and bring them in at their next appointment for scoring or they may complete the questionnaire during each scheduled appointment.
- 2. Add up ✓s by column. For every ✓: Several days = 1 More than half the days = 2 Nearly every day = 3
- 3. Add together column scores to get a TOTAL score.
- 4. Refer to the accompanying PHQ-9 Scoring Box to interpret the TOTAL score.
- 5. Results may be included in patient files to assist you in setting up a treatment goal, determining degree of response, as well as guiding treatment intervention.

Scoring: add up all checked boxes on PHQ-9

For every \checkmark Not at all = 0; Several days = 1; More than half the days = 2; Nearly every day = 3

Interpretation of Total Score

Total Score	Depression Severity		
1-4	Minimal depression		
 5-9	Mild depression		
 10-14	Moderate depression		
15-19	Moderately severe depression		
20-27	Severe depression		

PHQ9 Copyright © Pfizer Inc. All rights reserved. Reproduced with permission. PRIME-MD & is a trademark of Pfizer Inc.

A2662B 10-04-2005

CAGE Questionnaire

This is best used in a clinical setting as part of a general clinical history taking and may be phrased informally.

Have you ever felt you should Cut down on your drinking?	No Yes
Have people Annoyed you by criticizing your drinking?	No Yes
Have you ever felt bad or Guilty about your drinking?	No Yes
Have you ever had a drink first thing in the morning to steady your nerves or to get rid of a hangover (Eye opener)?	No Yes
$Total = \frac{1}{10} / 2$	1